

# Preventive Services Guide

Effective January 1, 2021

## **Ambetter Preventive Care Services**

Preventive care services can help you take charge of your health so you and your primary care provider (PCP) can catch problems before they start. These services include checkups, tests and screenings based on your age, weight or medical history.

See the charts on the following pages for the preventive services included in your Ambetter health plan. At your annual wellness exam, ask your PCP if you need any screenings or tests. Together, you and your PCP can stay updated about any changes in your health.

If you have any questions, talk to your doctor. Or you can call us at the toll-free number listed on the back of your Ambetter ID card.

#### **Ambetter Preventive Services Charts**

- 1. Adult Preventive Services
- 2. Women's Preventive Services
- 3. Children's Preventive Services

#### BENEFIT CONSIDERATIONS

Before using this guideline, please check your member specific benefit plan document and any federal or state mandates, if applicable. *Note: This is an overall guide to preventive care, but not all-inclusive.* 

Throughout this document the following acronyms are used:

- USPSTF: United States Preventive Services Task Force
- PPACA: Patient Protection and Affordable Care Act of 2010
- ACIP: Advisory Committee on Immunization Practices
- HHS: Health and Human Services
- HRSA: Health Resources and Services Administration

#### Ambetter's Preventive Services Guidelines

Preventive services include a broad range of benefits (including screening tests, counseling, and immunizations/vaccines). The federal Patient Protection and Affordable Care Act (PPACA) requires non-grandfathered health plans to cover preventive care services, when provided by network providers, without cost sharing to members. Preventive care services include: evidence based items or services that have in effect a rating of "A" or "B" in the current recommendations of the USPSTF, immunizations for routine use in children, adolescents and adults that have in effect a recommendation from the ACIP, with respect to infants, children and adolescents, evidence-informed preventive care and screenings provided for in the comprehensive guidelines supported by the HRSA and with respect to women, such additional preventive care and screenings as provided for in comprehensive guidelines supported by the HRSA.

To support your efforts and continuously improve the satisfaction of our members, we have adopted national practice parameters for disease management. Our goal in adopting national parameters is to help our members attain optimal quality of life. The parameters are provided to physicians for use as guidelines to assist them in clinical decision-making, and are not intended to be rigid standards.

#### **Adult Preventive Services**

**All members**: Annual wellness exams; all routine immunizations and vaccines recommended by the Advisory Committee on Immunization Practices of the Centers for Disease Control and Prevention (CDC).

**All members at an appropriate age and/or risk status:** Counseling and/or screening for: colorectal cancer; elevated cholesterol and lipids; certain sexually transmitted diseases; HIV; depression; high blood pressure; diabetes. Screening and counseling for alcohol abuse in a primary care setting; tobacco use; obesity; diet and nutrition.

**Men's health**: Intervention services as part of a full physical exam or periodic check-up for the purpose of education or counseling on potential health concerns, including smoking cessation counseling. Screening for prostate cancer for men age 40 and older; screening for abdominal aortic aneurysm in men 65-75 years old (USPSTF recommends this for males 65-75 years old who have smoked).

Routine Checkups <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Wellness Exam includes personal	Annually for ages 18-	oo oo jouro		00019000	oo jouro
history; blood pressure; body mass	21				mm alls
index (BMI); physical exam; preventive				A	nnually
screening; and counseling	Every 1–3 years, depending on risk factors				
Cancer Screenings <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Ĭ			<u>.                                      </u>		and women age 50-75 for
					ectal cancer
Colorectal Cancer Screening					est/fecal immunochemical
Colorectal Carreer Corectning					I DNA testing (Cologuard) y 3 years;
					y 5 years, oscopy every 5 years; or
					y every 10 years
	D	ationto at high righ for agl	aractal capacit due to family h	piotony or physical fact	oro.
	Patients at high risk for colorectal cancer due to family history or physical factors				UIS
Skin Cancer Screening		ms every 3 years at the healthcare provider	Annual total skin exar	m at discretion of your	healthcare provider
	discretion of your	· · · · · · · · · · · · · · · · · · ·	l nical breast exam and month	lv self-exam	
Breast Cancer Screening (Women)			vered once per year. High ris		-up mammograms may
		Ç	require an additiona		
Cervical Cancer Screening (Women)			21; if 30 years or older, either		
derried cancer careening (realism)			illomavirus (hrHPV) testing a		with hrHPV testing in
			65 years and older may stop		
Testicular and Prostate Cancer (Men)		Clinical testicular exam a	t each health maintenance v	isit and monthly self-e	xam
Other Recommended Screenings <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Body Mass Index (BMI)			healthcare provider in addition		
Body Maco Maco (Billi)	(can be scree	ened annually for overwei	ght and eating disorders, co	nsult the CDC's growtl	,
Abdominal Aartic Anguryem					Men between the ages of 65 to 75 that have ever
Abdominal Aortic Aneurysm					smoked
Blood Pressure (Hypertension)		At every acute/nonacut	e medical encounter and at I	east once every 2 yea	
Cholesterol Screening	E	very 5 years or more ofte	n at discretion of your health		
Diabetes Screening (Type 2)				s or earlier if risk facto	ers present
			Consider your risk factors, discuss with your		
			healthcare provider BMD		
Bone Mass Density (BMD) Test			testing for all post-	BMD test once, or m	ore often at the discretion
(Women)			menopausal women who	of your hea	lthcare provider
			have one or more risk		
			factors for osteoporosis fractures		
Hepatitis B Virus Infection Screening		Nonpregnant tee	ns and adults who have a hig	h risk for infection	
Infectious Disease Screening <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-con4 years	65+ years
Sexually Transmitted Infections (Chlamydia, Gonorrhea, Syphilis,	Annual screenings		ts under 25; annually for pati d under, if not previously vac		if at risk. HPV is for
and HPV 3) Tuberculosis screening: adults	Screenings re	ecommended for latent tu	berculosis infection in persor	ns who are at increase	d risk for infection
Immunizations 1, 2, 4, 5	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Immunizations 1, 2, 4, 5 Influenza Vaccine (Flu)	10-23 years	30-33 years	Annually	00-04 years	oor years
uozu vuoeo (r. u)	Δαρς 10+· Tda	n vaccines once (can sub	estitute 1-time dose for Td bo	oster) then boost with	Td every 10, years
Tetanus, Diphtheria, Pertussis (TD/Tdap)	•	t, talk to your doctor abou	it getting a Tdap vaccine dun aby from whooping cough (p	ing 3rd trimester of eve	, ,
Varicella Vaccine (Chicken Pox)	2 doses		no have not received the vac	<u> </u>	chicken pox
ζ	3 doses may be		exually active patients under 2		
	administered to both		under, if not previously vaccina		
Human Papillomavirus (HPV)	males and females		fit of HPV vaccination in this	0 0	
. ,	ages 19-26 with discretion from your	making is recomme	ended because some person	•	tely vaccinated might
	discretion from your benefit. healthcare provider				
Shingles Vaccine				50 yea	rs and older
Pneumococcal 13-Valent Conjugate (PCV13)		1 time dose	prior to age 65		1 dose 65> if no evidence of prior immunization
Pneumococcal Polysaccharide (PPSV23)	1 or 2 doses prior to age 65				1 dose 65> if no evidence of prior immunization
Meningococcal Vaccine	1 or more doses if not previously immunized, depending on risk factors and other indicator				
Hepatitis A Vaccine	2 doses if risk factors are present (if you did not get as a child)				
Hepatitis B Vaccine	3 doses if risk factors are present (if you did not get as a child)				
Haemophilus Influenza Type B (Hib)	(Pregnant women beginning at first prenatal visit. Consult with your healthcare provider)  1 or 3 doses if risk factors are present				
	1 or 2 doses for ac		pry of infection or previous		
Measles, Mumps, Rubella (MMR)	immunization				

#### **Women's Preventive Services**

Screenings for women's health, including: pregnancy-related preventive services, well-woman visits, preconception counseling and prenatal care, Pap tests and cervical cancer screening tests, contraceptive methods and counseling and screening and counseling for interpersonal and domestic violence.

Routine Checkups <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Wellness Exam includes personal history;	Annually for ages				ĺ
blood pressure; body mass index (BMI);	18-21			Annually	Annually
physical exam; preventive screening; and counseling	Every 1-3 years, depending on risk factors		Annually	Annually	
Routine Screenings <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Anemia Screening		Pregnan	t Women		
Cervical Cancer Screening (Women)	Initial pap test every 3 years beginning at age 21; if 30 years or older, either a Pap every 3 years alone or every 5 years screening with high-risk human papillomavirus (hrHPV) testing alone or every 5 years with hrHPV testing in combination with Pap (cotesting). Women 65 years and older may stop screening.				
FDA Approved Contraceptive Methods and Counseling	As prescribed by a healthcare provider for women with reproductive capability <sup>6</sup>				
Colorectal Cancer Screening	Screening for men and women age 50-75 for colorectal cancer Fecal occult blood test/fecal immunochemical test annually; or fecal DNA testing (Cologuard every 3 years; or flexible sigmoidoscopy every 5 years; or colonoscopy every 10 years				
	Patients	at high risk for colorecta	al cancer – 1 screening eve	ery 2 years	
Gestational Diabetes Screening			or those at high risk of dev		etes
Skin Cancer Screening	Periodic total skin exams every 3 years at discretion of your healthcare provider  Annual total skin exam at discretion of your healthcare provider			ılthcare provider	
Breast Cancer Screening	Annual clinical breast exam and monthly self-exam				
	Mammograms are covered once per year. High risk or necessary follow-up mammograms may require an additional annual visit.**				
Domestic and Interpersonal Violence Screening and Counseling	Recommended for all women with a routine screening and counseling by a network provider				
Breast Feeding and Post-Partum Counseling, Equipment and Supplies	For women as part of pre/post-natal counseling for pregnant women, with rental or purchase of certain breast feeding equipment through approved vendors				
Screening for Anxiety	Screening for anxiety in adolescent and adult women, including those pregnant or postpartum (clinical judgement should be used to determine screening frequency)			linical judgement	
Other Recommended Screenings <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Body Mass Index (BMI)			nealthcare provider in addi ht and eating disorders, c		
Blood Pressure (Hypertension)	At e	very acute/nonacute me	dical encounter and at lea	st once every 2 years	
Cholesterol Screening	At every acute/nonacute medical encounter and at least once every 2 years  Women ages 20 to 45 years for lipid disorders if at increased risk for coronary heart disease  Screenings every 5 years or more at age 45 and older as healthcare provider suggest				
Diabetes Screening (Type 2)	Every 3 years, beginning at age 45 or more often and beginning at younge age at the discretion of your healthcare provider				
Bone Mass Density (BMD) Test (Women)			Consider your risk factors, discuss with your healthcare provider. BMD testing for all postmenopausal women who have one or more risk factors for osteoporosis fractures	BMD test once, or m discretion of your he	
Infectious Disease Screening <sup>1</sup>	18-29 years	30-39 years	40-49 years	50-64 years	65+ years
Sexually Transmitted Infections (Chlamydia, Gonorrhea, Syphilis, and HPV 3)	Annual screenings for sexually active patients under 25; annually for patients age 25 and over if at risk.  HPV is for age 26 and under, if not previously vaccinated.				
Hepatitis B	3 doses if risk factors are present (if you did not get as a child) (Pregnant women beginning at first prenatal visit. Consult with your healthcare provider)			ovider)	

### **Children's Preventive Services**

Includes annual well child visits, screening newborns for hearing problems, thyroid disease, phenylketonuria, sickle cell anemia, and standard metabolic screening panel for inherited enzyme deficiency diseases. Counseling for fluoride for

prevention of dental cavities; screening for major depressive disorders; vision; lead; tuberculosis; developmental/autism; counseling for obesity.

Screening Tests <sup>1</sup>	0–1 year (Infancy)	1–4 years (Early Childhood)	5–11 years (Middle Childhood)	12–17 years (Adolescence)	
Well Baby Visits and Care (including cholesterol screening, height, weight, developmental milestones, and BMI)	Ages 1-2 weeks; and 1, 2, 4, 6, 9, and 12 months. Assess breastfeeding infants between 3–5 days of age	Ages 15, 18, and 24 months; and 3 and 4 years	Annually	Annually	
Anemia	Once between ages 9-12 months	As needed at the discretion of your healthcare provider		Starting at age 12, screen all non- pregnant adolescents for anemia every 5-10 years during well visit. Annually screen for anemia if at high risk	
Blood Test for Lead	Initial screening between ages 9-12 months	Annually at ages 2 and 3 years, and again at 4 years if in areas of high risk			
Urinalysis		Once at age 5 at the discretion of your healthcare provider			
Blood Pressure			Annually beginning at a	ge 3	
Hearing	Assess prior to discharge, or	Audiome	etry at ages 4, 5, 6, 8, 10,	12. 15. and 17	
Vision	by 1 month Assess prior to discharge, and by 6 months	• • • • • • • • • • • • • • • • • • • •		17 screen for strabismus (lazy eye)	
Pap Smear (Females)			Per ACS every 3 yrs. beginning at age 21 or as recommended by practitioner for abnormal findings		
Chlamydia screening				If sexually active and < 24	
Tests for Sexually Transmitted	Annual screenings for sexually active patients under 25; annually for patient				
Diseases		HPV is for age 26 and under	, if not previously vaccina		
Testicular Exam (Males)				Clinical exam and self-exam instruction annually beginning at age 15	
Congenital Hypothyroidism Screening	Newborns				
Critical Congenital Heart Disease Screening	Newborns before discharge from hospital				
Cholesterol/Lipid Disorders Screening	At-risk children 2-8 At-risk from 9 -11			At-risk adolescents 12-18	
Tuberculin Test		Children and ad	olescents at risk		
Routine Eye Exam for Children		1 visit a	nnually		
Depression	0.4	4.4	E 44 veers	Ages 11 - 17 12–17 years	
Immunizations <sup>1, 2, 4, 5</sup>	0–1 year (Infancy)	1–4 years (Early Childhood)	5–11 years (Middle Childhood)	(Adolescence)	
Hepatitis A	(intaney)			nd high-risk children over 24 months	
Hepatitis B	2 doses routinely recommended at birth and ages 1–2 months	1 doses 6–18 months	,		
Diphtheria, Tetanus, Pertussis (DTaP) Tetanus, Diphtheria, and Acellular Pertussis (Tdap) [Note: replaces Tetanus Diphtheria (Td)]	3 doses of DTaP routinely recommended at ages 2, 4, and 6 months	1 dose at 15–18 months	1 dose between 4–6 years	1 dose of Tdap between ages 7-10 instead of Td vaccine if you do not know if your child has received these; also between ages 13–18 years who missed Td booster at 11–12	
Polio Vaccine	2 doses routinely recommended at ages 2 and 4 months	1 dose recommended between 6–18 months	1 dose between 4–6 years		
Haemophilus (Hib)	3 doses routinely recommended at ages 2, 4, and 6 months	1 dose between 12–15 months			
Measles, Mumps, Rubella (MMR)		1 dose routinely recommended between 12–15 months	years		
Varicella Vaccine (Chicken Pox)		1 dose routinely recommended			
Pneumococcal Vaccine	3 doses routinely recommended at ages 2, 4, and 6 months	between 12–15 months  1 dose between 12–15 months	years		
Meningococcal Vaccine	5114 G 11011110	Certain high-risk group only. As needed at discretion of your healthcare provider		dose between ages 11–12 years; 1 dose at high school or college entry if     not previously vaccinated	

Human Papillomavirus (HPV)		3 doses between ages 11–12 years for males and females; Any dose not administered at the recommended age, should be administered at a subsequent visit	
Influenza Vaccine (Flu)	Annually for children 6 months of age and older		
Rotavirus	3 doses at 2, 4, and 6 months		

- 1 Ambetter will cover additional preventive benefits when required by the state.
- Some immunizations are indicated for certain conditions, discuss with your provider what routine preventive care and immunizations are best for you.
- 3 HPV is for age 26 and under if not previously vaccinated.
- <sup>4</sup> Ambetter covers vaccines under the preventive service benefit, without cost sharing, when services are rendered by an in-network provider and/or pharmacy who administers these vaccines.
- <sup>5</sup> Routine recommendation ask your primary care provider (PCP) about immunizations you may need.
- 6 Washington State FDA approved over-the-counter contraceptive devices and supplies are covered without a prescription.
- \*\*Ambetter pays for breast cancer screening once a year. When administered as a preventive breast imaging screening, digital breast tomosynthesis (known as 3-D mammography) is considered a covered, preventive benefit. *Note: Diagnostic mammograms are covered, but not part of preventive care coverage. Please work with your provider, for additional information.*

#### **Coverage Limitations and Exclusions**

- 1. Services not covered under the preventive care benefit may be covered under another portion of the medical benefit plan.
- 2. Generally, the cost of drugs, medications, vitamins, supplements, or over-the-counter items is not eligible as a preventive care benefit. However, certain outpatient prescription medications, tobacco cessation drugs and/or over the counter items, as required by PPACA, may be covered under the preventive benefit. For details, please refer to the member-specific pharmacy plan administrator.
- 3. An immunization is not covered if it does not meet company Vaccine Policy requirements for FDA labeling (including age and/or gender limitations) and if it does not have definitive ACIP recommendations published in the CDC's Morbidity and Mortality Weekly Report (MMWR).
- 4. Examinations, screenings, testing, or immunizations are not covered when:
  - a. required solely for the purposes of career, education, sports or camp, travel (including travel immunizations), employment, insurance, marriage or adoption, or
  - b. related to judicial or administrative proceedings or orders, or
  - c. conducted for purposes of medical research, or
  - d. required to obtain or maintain a license of any type.
- 5. Services that are investigational, experimental, unproven or not medically necessary are not covered. Please see applicable Medical Policies (EOC, SOB, etc.) for details.
- 6. Breastfeeding equipment and supplies not listed in the Indications for Coverage section above. This includes, but is not limited to:
  - a. Manual breast pumps and all related equipment and supplies.
  - b. Hospital-grade breast pumps and all related equipment and supplies.
  - c. Equipment and supplies not listed in the Covered Breastfeeding Equipment section above, including but not limited to:
    - i. Batteries, battery-powered adaptors, and battery packs.
    - ii. Electrical power adapters for travel.
    - iii. Bottles that are not specific to breast pump operation. This includes the associated bottle nipples, caps and lids.
    - iv. Travel bags, and other similar travel or carrying accessories.
    - v. Breast pump cleaning supplies including soap, sprays, wipes, steam cleaning bags and other similar products.
    - vi. Baby weight scales.
    - vii. Garments or other products that allow hands-free pump operation.
    - viii. Breast milk storage bags, ice-packs, labels, labeling lids, and other similar products.
    - ix. Nursing bras, bra pads, breast shells, nipple shields, and other similar products.
    - x. Creams, ointments, and other products that relieve breastfeeding related symptoms or conditions of the breasts or nipples.

The benefits within this document are currently effective unless otherwise noted. Always refer to your Schedule of Benefits and Evidence of Coverage to understand if there are any costs associated with your preventive care exam. It is important to know what type of service you are getting. If a non-preventive service is performed during the same healthcare visit/exam as a preventive service, you may have cost share (copayment, coinsurance and/or deductible charges) associated with the non-preventive care. If you receive any other covered services during a preventive care visit, you will be responsible to pay the applicable cost share for those other services.

#### ADDITIONAL PREVENTIVE SERVICES DETAILS

This Coverage Determination Guideline provides assistance in interpreting Ambetter preventive care services. When deciding coverage, the member specific benefit plan document must be referenced. This document is supplemental to your benefit plan document (e.g. Evidence of Coverage (EOC) and Schedule of Benefits (SOB), Member Handbook) and should not be used to guarantee coverage. Providers must first identify member eligibility, any federal or state regulatory requirements, and the member specific benefit plan coverage prior to use of this Coverage Determination Guideline. Other Policies and Coverage Determination Guidelines may apply; members should refer back to the EOC for detailed coverage information, including the essential health benefit plan. Ambetter reserves the right, in its sole discretion, to modify its Policies and Guidelines as necessary. This Coverage Determination Guideline is provided for informational purposes, your plan may not pay for all services and treatments in this guide. It does not constitute medical advice.

Note: Preventive services do not generally include services intended to treat an existing illness, injury, or condition. Benefits will be determined based on how the provider submits the bill. Claims must be submitted with the appropriate diagnosis and procedure code in order to be paid at the 100% benefit level. If during your preventive services visit you receive services to treat an existing illness, injury or condition, you may be required to pay a copay, deductible and/or coinsurance for those covered services.

This information is intended as a reference tool for your convenience and is not a guarantee of payment.

#### Statement of Non-Discrimination

Ambetter from Coordinated Care Corporation complies with applicable Federal and Washington state civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex, gender identity or sexual identity. Ambetter from Coordinated Care does not exclude people or treat them differently because of race, color, national origin, age, disability, or sex, gender identity or sexual orientation.

Ambetter from Coordinated Care:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
  - Qualified sign language interpreters
  - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
  - · Qualified interpreters
  - Information written in other languages

If you need these services, contact Ambetter from Coordinated Care at 1-877-687-1197 (TTY/TDD 1-877-941-9238).

If you believe that Ambetter from Coordinated Care has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, gender identity or sexual orientation, you can file a grievance with: Grievances Coordinator, Coordinated Care, 1145 Broadway, Suite 300, Tacoma, WA 98402, 1-877-687-1197 (TTY/TDD 1-877-941-9238), Fax 1-855-218-0588. You can file a grievance by mail, fax, or email <a href="https://www.way.uc.unit.com">www.way.uc.unit.com</a>. If you need help filing a grievance, Ambetter from Coordinated Care is available to help you. You can also file a civil rights complaint with:

- The U.S. Department of Health and Human Services, Office for Civil Rights electronically through the
  Office for Civil Rights Complaint Portal, available at https://ocrportal.hhs.gov/ocr/portal/lobby.jsf, or by
  mail or phone at: U.S. Department of Health and Human Services, 200 Independence Avenue SW.,
  Room 509F, HHH Building, Washington, DC 20201, 1-800-368-1019, 800-537-7697 (TDD). Complaint
  forms are available at http://www.hhs.gov/ocr/office/file/index.html.
- The Washington State Office of the Insurance Commissioner, electronically through the Office of the Insurance Commissioner Complaint portal available at https://www.insurance.wa.gov/file-complaint-orcheck-your-complaint-status, or by phone at 800-562-6900, 360-586-0241 (TDD). Complaint forms are available at https://fortress.wa.gov/oic/onlineservices/cc/pub/complaintinformation.aspx.



Onesalaka	Si usted, o alguien a quien está ayudando, tiene preguntas acerca de Ambetter de Coordinated Care Corporation, tiene derecho a
Spanish:	obtener ayuda e información en su idioma sin costo alguno. Para hablar con un intérprete, llame al 1-877-687-1197 (TTY/TDD 1-877-941-9238).
Chinese:	如果您·或是您正在協助的對象·有關於 Ambetter from Coordinated Care Corporation 方面的問題,您有權利免費以您的母語得到
Omnese.	幫助和訊息。如果要與一位翻譯員講話·請撥電話 1-877-687-1197 (TTY/TDD 1-877-941-9238)。
Vietnamese:	Nếu quý vị, hay người mà quý vị đang giúp đỡ, có câu hỏi về Ambetter from Coordinated Care Corporation, quý vị sẽ có quyền được giúp và có thêm thông tin bằng ngôn ngữ của mình miễn phí. Để nói chuyện với một thông dịch viên, xin gọi 1-877-687-1197 (TTY/TDD 1-877-941-9238).
Korean:	만약 귀하 또는 귀하가 돕고 있는 어떤 사람이 Ambetter from Coordinated Care Corporation 에 관해서 질문이 있다면 귀하는 그러한 도움과 정보를 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 그렇게 통역사와 얘기하기 위해서는 1-877-687-1197 (TTY/TDD 1-877-941-9238)로 전화하십시오.
Russian:	В случае возникновения у вас или у лица, которому вы помогаете, каких-либо вопросов о программе страхования Ambetter from Coordinated Care Corporation вы имеете право получить бесплатную помощь и информацию на своем родном языке. Чтобы поговорить с переводчиком, позвоните по телефону 1-877-687-1197 (TTY/TDD 1-877-941-9238).
Tagalog:	Kung ikaw, o ang iyong tinutulangan, ay may mga katanungan tungkol sa Ambetter from Coordinated Care Corporation, may karapatan ka na makakuha nang tulong at impormasyon sa iyong wika ng walang gastos. Upang makausap ang isang tagasalin, tumawag sa 1-877-687-1197 (TTY/TDD 1-877-941-9238).
Ukrainian:	В разі виникнення у вас або особи, якій ви допомагаєте, будь-яких запитань щодо програми страхування Ambetter from Coordinated Care Corporation ви маєте право отримати безкоштовну допомогу та інформацію на своїй рідній мові. Щоб поговорити з перекладачем, зателефонуйте за номером 1-877-687-1197 (TTY/TDD 1-877-941-9238).
Mon-Khmer, Cambodian:	ប្រសិនលោកអ្នកឬ នរណាម្នាក់ដែលអ្នកកំពុងតែដួយមានបញ្ហាអំពី Ambetter from Coordinated Care Corporation អ្នកមានសិទ្ធិទទួលបាន ជំនួយនិងព័ត៌មានជាភាសាលោកអ្នកដោយឥតគិតថ្លៃ។ សូមនិយាយទៅកាន់អ្នកបកប្រែតាមលេខ 1-877-687-1197 (TTY/TDD 1-877- 941-9238)
Japanese:	Ambetter from Coordinated Care Corporation について何かご質問がございましたらご連絡ください。 ご希望の言語によるサポートや情報を無料でご提供いたします。通訳が必要な場合は、1-877-687-1197 (TTY/TDD 1-877-941-9238) までお電話ください。
Amharic:	እርስዎ ወይም እርሰዎ የሚርዱት ሰው ስለ Ambetter from Coordinated Care Corporation ግብር ጥያቄ ካለዎት ያለምንም ወጪ በቋንቋዎ ድ <i>ጋ</i> ፍም እንዲሁም መረጃ የጣባፕት መብት አለዎት፤ ፤ አስተርጓሚ ለጣነ <i>ጋገ</i> ር በ 1-877-687-1197 (TTY/TDD 1-877-941-9238) ይደውሉ፤ ፤
Cushite:	Yoo sii ykn namaa gargaaraa jirtuu wa'ee Ambetter from Coordinated Care Corporation (Kuununsaa Qindeeffamaa) irra gaaffi qabaatan ta'ee gargaarsaa fi odeeffanoo afaan ketiin kaffaltii alla argachuuf mirgaa qabdaa. Turjumaana wajiin dubadhuu,1-877-687-1197 irra bilbilli (TTY/TDD 1-877-941-9238).
Arabic:	إذا كان لديك أو لدى شخص تساعده أسئلة حول Ambetter from Coordinated Care Corporation، لديك الحق في الحصول على المساعدة والمعلومات الضرورية بلغتك من دون أية تكلفة. للتحدث مع مترجم اتصل بـ 1197-687-871 (9238-941-877-1-787).
Punjabi:	ਜੇ ਤੁਹਾਡੇ, ਜਾਂ ਤੁਹਾਡੀ ਮਦਦ ਲੈ ਰਹੇ ਕਿਸੇ ਵਿਅਕਤੀ ਦੇ ਮਨ ਵਿਚ Ambetter from Coordinated Care Corporation ਦੇ ਬਾਰੇ ਕੋਈ ਸਵਾਲ ਹਨ. ਤਾਂ ਤੁਹਾਨੂੰ ਆਪਣੀ ਭਾਸ਼ਾ ਵਿਚ ਮੁਫਤ ਮਦਦ ਲੈਣ ਦਾ ਪੂਰਾ ਹੱਕ ਹੈ। ਦੁਭਾਸ਼ੀਏ ਨਾਲ ਗੱਲ ਕਰਨ ਲਈ 1-877-687-1197 (TTY/TDD 1-877-941-9238)'ਤੇ ਕਾਲ ਕਰੋ।
	Falls Sie oder jemand, dem Sie helfen, Fragen zu Ambetter from Coordinated Care Corporation hat, haben Sie das Recht,
German:	kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Um mit einem Dolmetscher zu sprechen, rufen Sie bitte die Nummer 1-877-687-1197 (TTY/TDD 1-877-941-9238) an.
Laotian:	ຖ້າທ່ານ ຫຼືຄົນທີ່ທ່ານກຳລັງຊ່ວຍເຫຼືອ ມີຄຳຖາມກ່ຽວກັບ Ambetter from Coordinated Care Corporation, ທ່ານມີສິດທີ່ຈະໄດ້ຮັບການຊ່ວຍເຫຼືອແລະຂໍ້ມູນ ຂ່າວສານທີ່ເປັນພາສາຂອງທ່ານ ໂດຍບໍ່ມີຄ່າໃຊ້ຈ່າຍ. ເພື່ອຈະເວົ້າກັບນາຍພາສາ ໃຫ້ໂທຫາ 1-877-687-1197 (TTY/TDD 1-877-941-9238).