

Clinical Policy: Caudal or Interlaminar Epidural Steroid Injections

Reference Number: CP.MP.164

Last Review Date: 04/18

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Epidural steroid injections have been used for pain control in patients with radiculopathy, spinal stenosis, and nonspecific low back pain, despite inconsistent results as well as heterogeneous populations and interventions in randomized trials. Epidural injections are performed utilizing three approaches in the lumbar spine: caudal, interlaminar, and transforaminal. Generally, candidates for epidural steroid injection are individuals who have acute radicular symptoms or neurogenic claudication unresponsive to traditional analgesics and rest, with significant impairment in activities of daily living.

Policy/Criteria

It is the policy of health plans affiliated with Centene Corporation® that invasive pain management procedures performed by a physician are medically necessary when *the relevant criteria are met and the patient receives only one procedure per visit, with or without radiographic guidance.*

- I. It is the policy of health plans affiliated with Centene Corporation® that caudal or interlaminar epidural steroid injections (ESIs) are **medically necessary** for the following indications:
 - A. *One caudal or interlaminar ESI for acute pain* management (pain lasting < 3 months) is considered **medically necessary** when all of the following are met:
 1. There is severe radicular pain that interferes substantially with ADLs;
 2. Severe pain persists after treatment with NSAID and/or opiate (both ≥ 3 days or contraindicated/not tolerated);
 3. The member cannot tolerate chiropractic or physical therapy and the injection is intended as a bridge to therapy.
 - B. *Initial ESI for chronic pain*, all of the following:
 1. One caudal or interlaminar ESI is requested at one level in the cervical, thoracic or lumbar region;
 2. Persistent radicular pain has been caused by spinal stenosis, disc herniation or degenerative changes in the vertebrae, as confirmed by physical exam and imaging;
 3. Pain interferes with activities of daily living, and has lasted for at least 3 months;
 4. The member has failed to respond to conservative therapy including all of the following:
 - a. ≥ 6 weeks chiropractic, physical therapy or prescribed home exercise program;
 - b. NSAID ≥ 3 weeks or NSAID contraindicated or not tolerated;
 - c. ≥ 6 weeks activity modification;
 5. The member is not currently being treated with full anticoagulation therapy. For patients on warfarin, INR (international normalized ratio) should be ≤ 1.4 prior to the procedure. Discontinuing anti-platelet therapy is a clinical decision balancing risks

CLINICAL POLICY

Caudal or Interlaminar Epidural Steroid Injections

and benefits of the procedure on therapy, versus the underlying medical condition if not treated appropriately;

6. Absence of systemic infection or local infection at the site of a planned injection.

C. *Second caudal or interlaminar ESI for chronic pain that **did not** improve from the first ESI, all of the following:*

1. One ESI is requested at one level in the cervical, thoracic or lumbar region;
2. At least 2 weeks have passed since the first ESI.

D. *Second or subsequent caudal or interlaminar ESI for chronic pain that **improved** from the first or second ESI, all of the following:*

3. Initial injection(s) led to $\geq 50\%$ relief for at least 2 months, associated with functional improvement;
4. At least 3 months have passed since the last ESI;
5. Less than 4 injections have been administered within 12 months;
6. Less than 12 months have elapsed since the initial injection at the level requested.

II. It is the policy of health plans affiliated with Centene Corporation that *A third or subsequent caudal or interlaminar ESI for chronic pain that **did not** improve from the first two ESIs is considered **not medically necessary** because effectiveness has not been established.*

III. It is the policy of health plans affiliated with Centene Corporation that *continuation of injections beyond 12 months or more than 4 therapeutic injections is considered **not medically necessary** because effectiveness and safety has not been established. When more definitive therapies cannot be tolerated or provided, consideration will be made on a case by case basis.*

IV. It is the policy of health plans affiliated with Centene Corporation that *caudal or interlaminar ESI for any other indication or location is considered **not medically necessary** because effectiveness has not been established.*

Background

There is much debate on the efficacy and medical necessity of multiple interventions for managing spinal pain. Epidural glucocorticoid injections have been used for pain control in patients with radiculopathy, spinal stenosis, and nonspecific low back pain despite inconsistent results as well as heterogeneous populations and interventions in randomized trials. Epidural injections are performed utilizing 3 approaches in the lumbar spine: caudal, interlaminar, and transforaminal. Generally, candidates for epidural steroid injection are individuals who have acute radicular symptoms or neurogenic claudication unresponsive to traditional analgesics and rest, with significant impairment in activities of daily living. Epidural steroid injections have been used in the treatment of spinal stenosis for many years, and no validated long-term outcomes have been reported to substantiate their use. However, significant improvement in pain scores, have been reported at 3 months.

Zhai et al¹ conducted a meta-analysis to assess the effects of various surgical and nonsurgical modalities, including epidural injections, used to treat lumbar disc herniation (LDH) or

CLINICAL POLICY

Caudal or Interlaminar Epidural Steroid Injections

radiculitis. A systematic literature search was conducted to identify RCTs which compared the effect of local anesthetic with or without steroids. The outcomes included pain relief, functional improvement, opioid intake, and therapeutic procedural characteristics. The reviewers concluded the meta-analysis confirms that epidural injections of local anesthetic with or without steroids have beneficial but similar effects in the treatment of patients with chronic low back and lower extremity pain.

Results of a 2 year follow-up of 3 randomized, double-blind, controlled trials, with a total of 360 patients with chronic persistent pain of disc herniation receiving either caudal, lumbar interlaminar or transforaminal epidural injections, showed similar efficacy of the 3 techniques with local anesthetic alone or local anesthetic with steroid. Caudal and interlaminar trials used in the assessment showed some superiority of steroids over local anesthetic, at 3 and 6 month follow-up. Interlaminar with steroids were superior to transforaminal at 12-months.²

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2018, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
62320	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; without imaging guidance
62321	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; with imaging guidance (ie, fluoroscopy or CT)
62322	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); without imaging guidance
62323	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); with imaging guidance (ie, fluoroscopy or CT)
62324	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (eg, anesthetic,

CLINICAL POLICY

Caudal or Interlaminar Epidural Steroid Injections

CPT® Codes	Description
	antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, cervical or thoracic; without imaging guidance
62325	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, cervical or thoracic; with imaging guidance (ie, fluoroscopy or CT)
62326	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); without imaging guidance
62327	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); with imaging guidance (ie, fluoroscopy or CT)

HCPCS Codes	Description
N/A	

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

+ Indicates a code requiring an additional character

ICD-10-CM Code	Description
M47.22	Other spondylosis with radiculopathy, cervical region
M47.23	Other spondylosis with radiculopathy, cervicothoracic region
M47.24	Other spondylosis with radiculopathy, thoracic region
M47.25	Other spondylosis with radiculopathy, thoracolumbar region
M47.26	Other spondylosis with radiculopathy, lumbar region
M47.27	Other spondylosis with radiculopathy, lumbosacral region
M48.00- M48.08	Spinal Stenosis
M50.10- M50.13	Cervical disc disorder with radiculopathy
M51.14- M51.17	Thoracic, thoracolumbar and lumbosacral intervertebral disc disorders with radiculopathy
M54.12	Radiculopathy, cervical region
M54.13	Radiculopathy, cervicothoracic region
M54.14	Radiculopathy, thoracic region
M54.15	Radiculopathy, thoracolumbar region
M54.16	Radiculopathy, lumbar region

CLINICAL POLICY
Caudal or Interlaminar Epidural Steroid Injections

ICD-10-CM Code	Description
M54.17	Radiculopathy, lumbosacral region
M54.5	Low back pain
M54.6	Pain in thoracic spine
M96.1	Postlaminectomy syndrome, not elsewhere classified

Reviews, Revisions, and Approvals	Date	Approval Date
Caudal and interlaminar ESI criteria reviewed in CP.MP.118	04/18	04/18
Split from CP.MP.118 Injections for Pain Management. No criteria changes.	08/18	

References

1. Zhai J, Zhang L, Li M, et al. Epidural injection with or without steroid in managing chronic low back and lower extremity pain: a meta-analysis of ten randomized controlled trials. *Int J Clin Exp Med*. 2015 Jun 15;8(6):8304-16. eCollection 2015.
2. Manchikanti L, Singh V, Pampati V, et al. Comparison of the efficacy of caudal, interlaminar, and transforaminal epidural injections in managing lumbar disc herniation: is one method superior to the other? *Korean J Pain*. 2015 Jan;28(1):11-21.
3. American College of Occupational and Environmental Medicine Guidelines: Chronic Pain. ACOEM 2008.
4. Chou R, Hashimoto R, Friedly J, Fu Rochelle, Dana T, Sullivan S, Bougatsos C, Jarvik J. Pain Management Injection Therapies for Low Back Pain. Technology Assessment Report ESIB0813. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. HHS 290-2012-00014-I.) Rockville, MD: Agency for Healthcare Research and Quality; March 2015.
5. Chou R. Subacute and chronic low back pain: Nonsurgical interventional treatment. In: *UpToDate, Atlas SJ (Ed), UpToDate, Waltham, MA*. Accessed 8/8/18.
6. Chou R, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. 2007;147:478-491.
7. Chou R et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain. An evidence-based clinical practice guideline from the American Pain Society. *Spine* 2009; 34: 1066-1077.
8. Chou R, Hashimoto R, Friedly J, et al. Epidural Corticosteroid Injections for Radiculopathy and Spinal Stenosis: A Systematic Review and Meta-analysis. *Annals Of Internal Medicine*, 163(5), 373-381. doi:10.7326/M15-0934
9. Heggeness MH. AAOS endorses back pain guidelines. *AAOS Now*. Sept 2010.
10. Manchikanti L et al. A Critical Review of the American Pain Society Clinical Practice Guidelines for Interventional Techniques: Part 1. Diagnostic Interventions. *Pain Physician* 2010; 13:E141-E174.
11. Manchikanti L et al. A Critical Review of the American Pain Society Clinical Practice Guidelines for Interventional Techniques: Part 2. Therapeutic Interventions. *Pain Physician* 2010; 13:E215-E264.

CLINICAL POLICY

Caudal or Interlaminar Epidural Steroid Injections

12. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in spinal pain. Part II: guidance and recommendations. *Pain Physician* 2013; 16: S49-S283.
13. Novak S, Nemeth WC. The basis for recommending repeating epidural steroid injections for radicular low back pain: a literature review. *Arch Phys Med Rehabil* 2008; 89:543-552.
14. Sharma AK, Vorobeychik Y, Wasserman R, et al. The Effectiveness and Risks of Fluoroscopically Guided Lumbar Interlaminar Epidural Steroid Injections: A Systematic Review with Comprehensive Analysis of the Published Data. *Pain Med.* 2016.
15. Staal JB et al. Injection therapy for subacute and chronic low-back pain. *Cochrane Database of Systematic Reviews* 2008, Issue 3. Art. No. CD001824. DOI: 10.1002/14651858.CD001824.pub3.
16. Vorobeychik Y, Sharma A, Smith CC, et al. The Effectiveness and Risks of Non-Image-Guided Lumbar Interlaminar Epidural Steroid Injections: A Systematic Review with Comprehensive Analysis of the Published Data. *Pain Med.* 2016.
17. Work Loss Data Institute. Low back – lumbar & thoracic (acute & chronic). Encinitas (CA): Work Loss Data Institute; 2011. Various p.
18. Hayes Medical Technology Directory. Epidural Steroid Injections for Low Back Pain and Sciatica. Jan 2013. Update Jan. 2017. Archived Mar. 2018.
19. Akuthota V, Bogduk N, Easa JE et al. Lumbar Transforaminal Epidural Steroid Injections: Review and Recommendation. *North American Pain Society.* Jan 2013. <https://www.spine.org/ResearchClinicalCare/QualityImprovement/ReviewRecommendations>
20. Smith CC, Booker T, Schaufele MK, Weiss P. Interlaminar versus transforaminal epidural steroid injections for the treatment of symptomatic lumbar spinal stenosis. *Pain Med.* 2010 Oct;11(10):1511-5.
21. Schaufele MK, Hatch L, Jones W. Interlaminar versus transforaminal epidural injections for the treatment of symptomatic lumbar intervertebral disc herniations. *Pain Physician.* 2006 Oct;9(4):361-6.
22. Chang-Chien GC, Knezevic NN, McCormick Z, et al. Transforaminal versus interlaminar approaches to epidural steroid injections: a systematic review of comparative studies for lumbosacral radicular pain. *Pain Physician.* 2014 Jul-Aug;17(4):E509-24.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

CLINICAL POLICY

Caudal or Interlaminar Epidural Steroid Injections

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

Note: For Medicare members, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international

CLINICAL POLICY**Caudal or Interlaminar Epidural Steroid Injections**

copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.