

Clinical Policy: Selective Nerve Root Blocks and Transforaminal Epidural Steroid Injections

Reference Number: CP.MP.165

Last Review Date: 04/18

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Transforaminal epidural steroid injections (TFESIs) and selective nerve route blocks (SNRBs) are alternatives to interlaminar epidural steroid injections for the treatment of radicular pain. SNRBs consist of a small amount of local anesthetic injected adjacent to a spinal nerve root, and are most often used to diagnose the source of pain.¹ During a TFESI, a larger amount of local anesthetic or corticosteroid is injected into the intervertebral foramen, where the injectate spreads to target multiple nerves. SNRBs and TFESIs share similar safety considerations, procedural techniques, and anatomical benchmarks.¹

Policy/Criteria

It is the policy of health plans affiliated with Centene Corporation® that invasive pain management procedures performed by a physician are **medically necessary** when *the relevant criteria are met and the patient receives only one procedure per visit, with or without radiographic guidance.*

I. Selective Nerve Root Blocks

A. *One selective nerve root block (SNRB) for chronic pain* is considered **medically necessary** to establish a diagnosis and confirm beneficial response when all the following criteria are met:

1. Request is for an SNRB with a local anesthetic at a single nerve root;
2. There is persistent radicular pain in a defined nerve root level and the diagnosis remains uncertain after standard evaluation (neurologic examination, radiological studies and electrodiagnostic studies);
3. Pain interferes with activities of daily living (ADLs) and has lasted for at least 3 months;
4. The member has failed to respond to conservative therapy including all of the following:
 - a. ≥ 6 weeks chiropractic, physical therapy or prescribed home exercise program;
 - b. Nonsteroidal anti-inflammatory drugs (NSAID) ≥ 3 weeks or NSAID contraindicated or not tolerated;
 - c. ≥ 6 weeks activity modification;
5. The member is not currently being treated with full anticoagulation therapy. For patients on warfarin, international normalized ratio (INR) should be ≤ 1.4 prior to the procedure.-
Discontinuing anti-platelet therapy is a clinical decision balancing risks and benefits of the procedure on therapy, versus the underlying medical condition if not treated appropriately;
6. Absence of systemic infection or local infection at the site of a planned injection.

- B.** *A second SNRB for chronic pain* is considered **medically necessary** when multilevel pathology is suspected and it has been at least two weeks since the prior injection.
- C.** *One SNRB for acute pain management* is considered **medically necessary** when all of the following are met:
 - 1. Pain has lasted for < 3 months;
 - 2. There is severe radicular pain in a specific nerve root distribution that interferes substantially with ADLs;
 - 3. Severe pain persists after treatment with NSAID and/or opiate (both ≥ 3 days or contraindicated/not tolerated);
 - 4. The member cannot tolerate chiropractic or physical therapy and the injection is intended as a bridge to therapy.
- D.** *SNRBs* are considered **not medically necessary** for any other indication because effectiveness has not been established.

II. Transforaminal Epidural Steroid Injections

- A.** *One transforaminal epidural steroid injection (TFESI) for chronic pain* in the lumbar region is considered **medically necessary** when all of the following are met:
 - 1. TFESI is requested for a single level bilaterally or two levels unilaterally;
 - 2. There is persistent radicular pain caused by disc herniation in a defined nerve root level, or spinal stenosis confirmed by physical exam and imaging;
 - 3. Pain interferes with ADLs and has lasted for at least 3 months;
 - 4. The member has failed to respond to conservative therapy including all of the following:
 - a. ≥ 6 weeks chiropractic, physical therapy or prescribed home exercise program;
 - b. NSAID ≥ 3 weeks or NSAID contraindicated or not tolerated;
 - c. ≥ 6 weeks activity modification;
 - 5. The member is not currently being treated with full anticoagulation therapy. For patients on warfarin, INR should be ≤ 1.4 prior to the procedure. Discontinuing anti-platelet therapy is a clinical decision balancing risks and benefits of the procedure on therapy, versus the underlying medical condition if not treated appropriately;
 - 6. Absence of systemic infection or local infection at the site of a planned injection.
- B.** *A second TFESI for chronic pain* in the lumbar region is considered **medically necessary** when meeting all of the following:
 - 1. There was not a positive response to the initial injection;
 - 2. Request is for a TFESI at a single level bilaterally or two levels unilaterally;
 - 3. At least two weeks have passed since the initial TFESI.
- C.** *Subsequent TFESIs for chronic pain* are considered **medically necessary** if symptoms recur after a favorable response to the diagnostic SNRB(s) or initial TFESI(s) and all of the following:
 - 1. The TFESI is requested at a single level bilaterally or two levels unilaterally;

2. There was $\geq 50\%$ relief associated with functional improvement for at least 2 months from the prior injection(s);
 3. TFESIs are given at least 3 months apart;
 4. Less than 4 injections have been given at the same site within 12 months;
 5. The member is not currently being treated with full anticoagulation therapy. For patients on warfarin, INR should be ≤ 1.4 prior to the procedure. Discontinuing anti-platelet therapy is a clinical decision balancing risks and benefits of the procedure on therapy, versus the underlying medical condition if not treated appropriately;
 6. Absence of systemic infection or local infection at the site of a planned injection.
- D.** *Continuation of injections beyond 12 months or more than 4 therapeutic injections* is considered **not medically necessary** because effectiveness and safety has not been established. When more definitive therapies cannot be tolerated or provided, consideration will be made on a case by case basis.
- E.** *One TFESI for acute pain management* is considered **medically necessary** when all of the following are met:
1. Pain has lasted for < 3 months;
 2. There is severe radicular pain in a specific nerve root distribution that interferes substantially with ADLs;
 3. Severe pain persists after treatment with NSAID and/or opiate (both ≥ 3 days or contraindicated/not tolerated);
 4. The member cannot tolerate chiropractic or physical therapy and the injection is intended as a bridge to therapy.
- F.** *TFESIs for any other indication or location* are considered **not medically necessary** because effectiveness has not been established.

Background

Epidural steroid injections/selective nerve root blocks

There is great controversy regarding the effectiveness of invasive interventions for spinal pain. Epidural glucocorticoid injections have been used for pain control in patients with radiculopathy, spinal stenosis, and nonspecific low back pain despite inconsistent results as well as heterogeneous populations and interventions in randomized trials. Epidural injections are performed utilizing 3 approaches in the lumbar spine: caudal, interlaminar, and transforaminal. Generally, candidates for epidural steroid injection are individuals who have acute radicular symptoms or neurogenic claudication unresponsive to traditional analgesics and rest, with significant impairment in activities of daily living. Epidural steroid injections have been used in the treatment of spinal stenosis for many years, and no validated long-term outcomes have been reported to substantiate their use. However, significant improvement in pain scores, have been reported at 3 months. A SNRB is primarily used to diagnose the specific source of nerve root pain. In a SNRB, a local anesthetic is used. When used for therapeutic indications, a steroid is added and it is usually referred to as a selective transforaminal epidural steroid injection.

Zhai et al conducted a meta-analysis to assess the effects of various surgical and nonsurgical modalities, including epidural injections, used to treat lumbar disc herniation (LDH) or radiculitis.² A systematic literature search was conducted to identify RCTs which compared the effect of local anesthetic with or without steroids. The outcomes included pain relief, functional improvement, opioid intake, and therapeutic procedural characteristics. The reviewers concluded that the meta-analysis confirms that epidural injections of local anesthetic with or without steroids have beneficial but similar effects in the treatment of patients with chronic low back and lower extremity pain.²

Results of a 2 year follow-up of 3 randomized, double-blind, controlled trials, with a total of 360 patients with chronic persistent pain of disc herniation receiving either caudal, lumbar interlaminar or transforaminal epidural injections, showed similar efficacy of the 3 techniques with local anesthetic alone or local anesthetic with steroid.³ Interlaminar injections with steroids were superior to transforaminal at 12-months.³

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2018, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
64479	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); cervical or thoracic, single level
64480	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); cervical or thoracic, each additional level (List separately in addition to code for primary procedure)
64483	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); lumbar or sacral, single level
64484	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); lumbar or sacral, each additional level (List separately in addition to code for primary procedure)

HCPCS Codes	Description
N/A	

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

+ Indicates a code requiring an additional character

ICD-10-CM Code	Description
G56.00-G56.93	Mononeuropathies of upper limb
G57.00-G57.93	Mononeuropathies of lower limb
M48.061-M48.062	Spinal stenosis, lumbar region
M50.00-M50.93	Cervical disc disorders
M51.04-M50.06	Thoracic, thoracolumbar, and lumbosacral intervertebral disc disorders with myelopathy
M51.14-M51.27	Thoracic, thoracolumbar and lumbosacral intervertebral disc disorders with radiculopathy
M54.12	Radiculopathy, cervical region
M54.13	Radiculopathy, cervicothoracic region
M54.14	Radiculopathy, thoracic region
M54.15	Radiculopathy, thoracolumbar region
M54.16	Radiculopathy, lumbar region
M54.17	Radiculopathy, lumbosacral region
M54.30-M54.32	Sciatica
M54.40-M54.42	Lumbago with sciatica
M54.5	Low back pain

Reviews, Revisions, and Approvals	Date	Approval Date
SNRB and TFESI criteria reviewed and updated in CP.MP.118.	04/18	04/18
TFESI criteria reviewed and updated in CP.MP.118 with criteria added for initial injections.	05/18	05/18
Split from CP.MP.118 Injections for Pain Management. Minor wording changes to criteria with no clinical impact. Background and references updated.	08/18	

References

1. Bicket MC, Benzon HT, Cohen SP. Essentials of pain medicine (Fourth Edition). Chapter 63 – Transforaminal Epidural Steroid Injections and Selective Nerve Root Blocks. Elsevier. 2018, p. 573-584.e2
2. Zhai J, Zhang L, Li M, et al. Epidural injection with or without steroid in managing chronic low back and lower extremity pain: a meta-analysis of ten randomized controlled trials. Int J Clin Exp Med. 2015 Jun 15;8(6):8304-16.
3. Manchikanti L, Singh V, Pampati V, et al. Comparison of the efficacy of caudal, interlaminar, and transforaminal epidural injections in managing lumbar disc herniation: is one method superior to the other? Korean J Pain. 2015 Jan;28(1):11-21.
4. American College of Occupational and Environmental Medicine Guidelines: Chronic Pain. ACOEM 2008.
5. Chou R, Hashimoto R, Friedly J, Fu Rochelle, Dana T, Sullivan S, Bougatsos C, Jarvik J. Pain Management Injection Therapies for Low Back Pain. Technology Assessment Report ESIB0813. (Prepared by the Pacific Northwest Evidence-based Practice Center under

- Contract No. HHS-2012-00014-I.) Rockville, MD: Agency for Healthcare Research and Quality; March 2015.
6. Chou R. Subacute and chronic low back pain: Nonsurgical interventional treatment. In: UpToDate, Atlas SJ (Ed), UpToDate, Waltham, MA. Accessed 8/8/18.
 7. Chou R, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med.* 2007;147:478-491.
 8. Chou R et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain. An evidence-based clinical practice guideline from the American Pain Society. *Spine* 2009; 34: 1066-1077.
 9. Chou R, Hashimoto R, Friedly J, et al. Epidural Corticosteroid Injections for Radiculopathy and Spinal Stenosis: A Systematic Review and Meta-analysis. *Annals Of Internal Medicine,* 163(5), 373-381. doi:10.7326/M15-0934
 10. Heggeness MH. AAOS endorses back pain guidelines. *AAOS Now.* Sept 2010.
 11. Manchikanti L et al. A Critical Review of the American Pain Society Clinical Practice Guidelines for Interventional Techniques: Part 1. Diagnostic Interventions. *Pain Physician* 2010; 13:E141-E174.
 12. Manchikanti L et al. A Critical Review of the American Pain Society Clinical Practice Guidelines for Interventional Techniques: Part 2. Therapeutic Interventions. *Pain Physician* 2010; 13:E215-E264.
 13. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in spinal pain. Part II: guidance and recommendations. *Pain Physician* 2013; 16: S49-S283.
 14. Novak S, Nemeth WC. The basis for recommending repeating epidural steroid injections for radicular low back pain: a literature review. *Arch Phys Med Rehabil* 2008; 89:543-552.
 15. Staal JB et al. Injection therapy for subacute and chronic low-back pain. *Cochrane Database of Systematic Reviews* 2008, Issue 3. Art. No. CD001824. DOI: 10.1002/14651858.CD001824.pub3.
 16. Vorobeychik Y, Sharma A, Smith CC, et al. The Effectiveness and Risks of Non-Image-Guided Lumbar Interlaminar Epidural Steroid Injections: A Systematic Review with Comprehensive Analysis of the Published Data. *Pain Med.* 2016.
 17. Work Loss Data Institute. Low back – lumbar & thoracic (acute & chronic). Encinitas (CA): Work Loss Data Institute; 2011. Various p.
 18. Hayes Medical Technology Directory. Epidural Steroid Injections for Low Back Pain and Sciatica. Jan 2013. Update Jan. 2017. Archived Mar. 2018.
 19. Akuthota V, Bogduk N, Easa JE et al. Lumbar Transforaminal Epidural Steroid Injections: Review and Recommendation. North American Pain Society. Jan 2013.
<https://www.spine.org/ResearchClinicalCare/QualityImprovement/ReviewRecommendations/statements>
 20. Singh JR, Cardozo E, Christolias GC. The Clinical Efficacy for Two-Level Transforaminal Epidural Steroid Injections. *PM R.* 2017 Apr;9(4):377-382.
 21. Smith CC, Booker T, Schaufele MK, Weiss P. Interlaminar versus transforaminal epidural steroid injections for the treatment of symptomatic lumbar spinal stenosis. *Pain Med.* 2010 Oct;11(10):1511-5.

22. Schaufele MK, Hatch L, Jones W. Interlaminar versus transforaminal epidural injections for the treatment of symptomatic lumbar intervertebral disc herniations. *Pain Physician*. 2006 Oct;9(4):361-6.
23. Chang-Chien GC, Knezevic NN, McCormick Z, et al. Transforaminal versus interlaminar approaches to epidural steroid injections: a systematic review of comparative studies for lumbosacral radicular pain. *Pain Physician*. 2014 Jul-Aug;17(4):E509-24.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

Note: For Medicare members, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.